

Name: \_\_\_\_\_  
 Period: \_\_\_\_\_

**Newton's Second Law and Weight**

*You already know that if you push harder on an object it will accelerate more. Also, if you push with the same force on a heavier object it will have less acceleration. You instinctive know Newton's Second Law of Motion.*


**Newton's Second Law:  $F = ma$**

The acceleration of an object is proportional to the force acting on it and inversely proportional to its mass.


OR

*More force causes more acceleration; more mass causes less acceleration.*

Same  $m$ ; smaller  $F$ ; smaller  $a$   
 $a = 20/10 = 2 \text{ m/s}^2 \rightarrow$




Same  $m$ ; bigger  $F$ ; bigger  $a$   
 $a = 40/10 = 4 \text{ m/s}^2 \rightarrow$




*More force causes more acceleration. Doubling the force, doubles the acceleration.*

Same  $F$ ; smaller  $m$ ; bigger  $a$   
 $a = 40/10 = 4 \text{ m/s}^2 \rightarrow$



Same  $m$ ; bigger  $F$ ; bigger  $a$   
 $a = 20/20 = 1 \text{ m/s}^2 \rightarrow$



*More mass causes less acceleration. Doubling the mass, halves the acceleration.*

**$F = ma$**

Force (in Newtons)  $\rightarrow$   $F = ma$   $\leftarrow$  Mass (in kg)

$\leftarrow$  Acceleration (in  $\text{m/sec}^2$ )

*Force equals mass times acceleration.*

*Ex. How big a force does it take to give a 50 kg object an acceleration of 40  $\text{m/s}^2$ .*

Variables: $m = 50 \text{ kg}$ $a = 40 \text{ m/s}^2$ $F = \underline{\hspace{2cm}}$	Equation: $F = ma$ Solve: $F = 50(40)$ $F = \underline{2000\text{N}}$
---	---

The acceleration of a mass is actually due to the **net force** acting on the object:

$F_{\text{net}} = \Sigma F = ma$

*You must add up all of the forces to find the acceleration.*


**Mass vs. Weight**

In everyday speech we use mass and weight interchangeably, but in science mass and weight are very different.

**Mass** is the amount matter in an object (all of its atoms and molecules).

**Weight** is the force of gravity pulling on mass.

*On Earth*




Mass  $\rightarrow m = 50 \text{ kg}$

Weight  $\rightarrow F = 500 \text{ N}$

*Mass is measured in kg;  
Weight is a force measured in N.*

*In Space*



Mass  $\rightarrow m = 50 \text{ kg}$

Weight = 0 N


*In space an object still has mass, because it still has its atoms and molecules, but there is no weight, because there is no gravity.*

**Acceleration Due to Gravity**

All objects fall with the same acceleration:  $-9.8 \text{ m/s}^2$ . This is known as the acceleration due to gravity ( $g$ ). *This is not the force of gravity.*

**$g$  is a constant:  $-9.8 \text{ m/s}^2$**

**The acceleration of all falling objects is the same on the earth.**




All objects fall towards the earth at  $-9.8 \text{ m/s}^2$ .

Each second it will be going 9.8  $\text{m/s}$  faster.

$a = -9.8 \text{ m/s}^2$  due to gravity

**Heavy and light object fall at the same rate.**



*If it wasn't for air friction a feather and a hammer would fall at the same rate:  $-9.8 \text{ m/s}^2$ . In air, the hammer's extra weight allows it to push thru the air molecules faster.*

**Measuring Weight**

If you already know mass it is easy to calculate weight: just multiply mass times  $9.8 \text{ m/s}^2$ .

*To make calculations easier we often use  $g = 10 \text{ m/s}^2$ .*

**$F_w = mg$**

Force of Weight (in Newtons)  $\rightarrow$   $F_w = mg$   $\leftarrow$  Mass (in kg)

$\leftarrow$  Acceleration due to gravity ( $9.8 \text{ m/sec}^2$ )




*Weight equals mass times the acceleration due to gravity.*

*Ex. Find the weight of a 2 kg mass.*

Variables: $m = 2 \text{ kg}$ $g = 10 \text{ m/s}^2$ $F_w = \underline{\hspace{2cm}}$	Equation: $F_w = mg$ Solve: $F = (2)(10)$ $F = 20 \text{ N}$
--	--

*Ex. Find the mass of a 2 N object.*

Variables: $F_w = 2 \text{ N}$ $g = 10 \text{ m/s}^2$ $m = \underline{\hspace{2cm}}$	Equation: $F_w = mg$ Solve: $2 = (m)(10)$ $m = 2/10 = 0.2 \text{ kg}$
---	---

1. Weight 2. Mass 3. N 4. $F = ma$ 5. g	A. The units of weight and force. B. Newton's Second Law mathematically. C. The acceleration due to gravity = $-9.8\text{m/s}^2$ . D. The force of gravity on matter. E. How much matter an object contains.	Which of Newton's Three Laws Applies: Law 1, 2, or 3?  <input type="checkbox"/> Pushing a cart down the hall, when you try to turn it it tries to go straight. <input type="checkbox"/> More acceleration takes more force. <input type="checkbox"/> When you push your knuckles into a table, it hurts your knuckles. <input type="checkbox"/> A ball thrown into the ground bounces back up.
<i>Will an object accelerate faster or slower?</i> If its mass is increased? If the force pulling on it decreases? If the force pushing on it increases? If its mass is decreased?		A 6 kg object experiences a $5\text{ m/s}^2$ acceleration? Find the force that caused this acceleration.  A 3 kg rock accelerates to the left at $12\text{ m/s}^2$ . Find the net force that caused this.  A 12 kg box is pushed to the left by a 48 N force. Find its acceleration.
Why is $F = ma$ not entirely correct?		 <p>A) Calculate the object's net force. B) Calculate the object's acceleration.</p>
<i>More, less, or the same as on the Earth?</i> When an astronaut lands on the moon: The astronaut's mass is: _____ The astronaut's weight is: _____ The astronaut's inertia is: _____		 <p>Calculate the object's acceleration.</p>
Without air friction, which falls faster, heavy or light objects? Why?  If there is air friction, which falls faster? Why?		Using the weight equation, calculate the weight of a 45 kg rock?
How fast is the acceleration due to gravity? If an object falls from rest, how fast will it be going: after 1 second? after 2 seconds? after 6 seconds?		 <p><i>Find the force pulling left.</i></p>
Calculate the mass of a 10 N apple.  What is the mass of a 100 gram apple in kilograms?  What is the weight of the above apple?  What is the weight of a 250 N object?  If 100 kg person weighed 400 N on the planet Zorg, what is the acceleration due to gravity on Zorg?		12 N pulls to the left and 20 N pulls to the right on a 2 kg object. Draw the problem:  Calculate the object's acceleration.