


**Speed vs. Velocity**

Velocity is speed with direction.

Example: A person walks 4 m/s—speed (no direction).



**Velocity changes when direction changes.**

**Scalars vs. Vectors**

Vectors require direction;  
 Scalars only need magnitude (how big).

Remember: Speed is a Scalar; Velocity is a Vector.

Vectors require magnitude (how much) and direction, often vectors can cancel each other out (not acceleration, though).

**12 m/s west**      Speed: 12 m/s.  
 Magnitude    Direction    Velocity: 12 m/s west.

**Acceleration**

Acceleration is how fast you change velocity OR how much the velocity changed in a certain amount of time.

An object accelerates when it changes speed OR changes direction!

$$\text{Acceleration (in m/s}^2\text{)} \rightarrow a = \frac{\Delta V}{\Delta T}$$

← Change of Velocity (in meters/sec)  
 ← Change of Time (in seconds)

Acceleration equal change of velocity divided by change of time.

$$\Delta V = V_{\text{final}} - V_{\text{initial}}, \text{ so, } a = \frac{V_{\text{final}} - V_{\text{initial}}}{\Delta T}$$

**Finding  $\Delta V$ .**

$\Delta$  always = final – initial.  
 $\Delta V = V_{\text{final}} - V_{\text{initial}}$  OR  
 Final velocity – Initial velocity.

If  $\Delta V$  is positive the object is speeding up.

If  $\Delta V$  is negative the object is slowing down (see below).

Ex. A plane starts at rest and ends up going 200 m/s in 10 secs. Calculate its acceleration.

**Step 1: Variables**  
 $V_i = 0 \text{ m/s}$  (at rest)  
 $V_f = 200 \text{ m/s}$   
 $T = 10 \text{ sec}$   
 $a = \underline{\hspace{2cm}}$

**Step 3: Put in numbers and solve**  
 $a = \frac{\Delta V}{\Delta T} = \frac{V_f - V_i}{\Delta T} = \frac{200 - 0}{10}$   
 $a = \frac{200}{10} = 20$

**Step 2: Formula**  
 $a = \frac{\Delta V}{\Delta T}$

**Step 4: Add units**      Pos. means speeding up  
 $a = 20 \text{ m/s}^2$

Ex. A race car starts at 40 m/s slows to 10 m/s in 5 seconds. Calculate the car's acceleration.

**Step 1: Variables**  
 $V_i = 40 \text{ m/s}$   
 $V_f = 10 \text{ m/s}$   
 $T = 5 \text{ sec}$   
 $a = \underline{\hspace{2cm}}$

**Step 3: Put in numbers and solve**  
 $a = \frac{\Delta V}{\Delta T} = \frac{V_f - V_i}{\Delta T} = \frac{10 - 40}{5}$   
 $a = \frac{-30}{5} = -6$

**Step 2: Formula**  
 $a = \frac{\Delta V}{\Delta T}$

**Step 4: Add units**      Neg. means slowing down  
 $a = -6 \text{ m/s}^2$

Negative acceleration means an object is slowing down OR speeding up in the negative direction. Slowing down is also called "deceleration".

**Distance and Acceleration**

An object that is accelerating will travel farther each second.

**Constant Speed—Equal Distance**



Points are equal distance, so velocity is constant. Since the velocity is constant, the initial and final velocity are equal and the acceleration equals zero.

**Positive Acceleration—Increasing Distance**



The distance between the points is increasing, so velocity is increasing. The object is accelerating: traveling faster each second and covering more distance every second.

**Measuring Acceleration**

To measure an object's acceleration you need to measure the object's velocity before and after the acceleration.

If the object starts at rest you know that  $V_i = 0 \text{ m/s}$ .  
 If the object stops you know that  $V_f = 0 \text{ m/s}$ .

Measure  $V_i$   
(Initial Velocity)

$V_i = \frac{\Delta D}{\Delta T} = \frac{4 \text{ m}}{1 \text{ sec}}$   
 $V_{\text{initial}} = 4 \text{ m/s}$

Measure  $\Delta T$   
(Time it took to Accelerate)

Accelerates for 2 seconds  
So  $\Delta T = 2 \text{ sec}$

$$a = \frac{V_f - V_i}{\Delta T} = \frac{8 - 4}{2}$$

$$V_{\text{initial}} = \frac{4}{2} = 2 \text{ m/s}^2$$


Measure  $V_f$   
(Final Velocity)


$V_f = \frac{\Delta D}{\Delta T} = \frac{8 \text{ m}}{1 \text{ sec}}$   
 $V_{\text{final}} = 8 \text{ m/s}$


Speed (S) or Velocity (V)	Scalar (S) or Vector (V)	Mass, Time, Distance, Velocity, or Acceleration?
<input type="checkbox"/> A bike goes 25 m/s toward main street. <input type="checkbox"/> A person walks 4 mph. <input type="checkbox"/> A plane flies 200 m/s. <input type="checkbox"/> A bird flies 100 mph due south.	<input type="checkbox"/> 40 mph toward Dallas. <input type="checkbox"/> 3 m/s <sup>2</sup> to the left. <input type="checkbox"/> 10 meters up the hill. <input type="checkbox"/> 12 meter per sec <sup>2</sup> . <input type="checkbox"/> Direction matters. <input type="checkbox"/> No direction is needed	<input type="checkbox"/> 2 hrs <input type="checkbox"/> 5 sec <input type="checkbox"/> 8 kg <input type="checkbox"/> 3 m/s <input type="checkbox"/> 9 mph <input type="checkbox"/> 4 m/s <sup>2</sup> <input type="checkbox"/> 6 mph/sec <input type="checkbox"/> 12 m <input type="checkbox"/> 1 in


*Accelerating? Yes, No, or Maybe?*


At constant velocity.  
 Going 5 m/s then going 3 m/s.  
 A car going around a corner. (see graphic at right).  
 At constant speed.  
 Stopping.  
 A car at rest.



Object A 

Object B 

Object C 

Object D 

*Choose which of the above applies to the following*

Constant speed.                     Distance increases  
 Positive acceleration.             Starts at rest.  
 At constant velocity.             Is stopping.  
 Accelerating.                     Constant direction.  
 Decelerating.                     Negative acceleration.  
 Acceleration = 0.                 V<sub>i</sub> = V<sub>f</sub>

*Object A accelerates at 10 m/s<sup>2</sup>; Object B accelerates at 5 m/s<sup>2</sup>.*

Which one will go faster?  
 Which one will take more time to reach a high speed?  
 If they start at rest, which one will reach 40 m/s first?  
 Which one goes farther (longer distance)?  
 Which one will be 100m away sooner?

*Give what you know for the following: (V<sub>i</sub>, V<sub>f</sub> or a)*

An object at constant velocity.  
 An object that is stopping.  
 An object that accelerates from rest.  
 An object at rest.

A person starts running from 2 m/s to 6 m/s in 2 seconds. Calculate the person's acceleration.

A dragster's top acceleration is 60 m/s<sup>2</sup>. If it starts from rest at the starting line, how fast will it be going after 3 seconds?

Variables:	Solve:
Formula:	

Variables:	Solve:
Formula:	

A plane stops from 250 mph in 25 seconds. Calculate the planes acceleration.

A car travels 30 m in 5 seconds. After accelerating for 3 seconds, it travels 20 m in 2 seconds. Calculate the car's acceleration.

Variables:	Solve:
Formula:	

- 1) Find V<sub>i</sub>.
- 2) Find V<sub>f</sub>.
- 3) Calculate a.